

Institution Name and Address:

**DIABETES MEDICAL MANAGEMENT PLAN  
INTENSIVE THERAPY**

Patient Label or MRN, Acct#, Patient Name, DOB, Date of Service

Page 1 of 3

**Part 2: Virginia Diabetes Medical Management Plan (DMMP)**

To be completed by physician/provider.

Notice to Parents: Medication(s) **MUST** be brought to school by the PARENT/GUARDIAN in a container that is appropriately labeled by the pharmacy or physician/practitioner.

In order for schools to safely administer medication during school hours, the following regulations should be observed:

- A new copy of the DMMP must be completed at the beginning of each school year. This form, an Authorization for Medication Administration form, or MD prescription must be received in order to change diabetes care at school during the school year.

<b>Student Name (Last, First, MI)</b>	Student's Date of Birth	
School	Student's Grade □	Home Phone
Parent Name	Work/Cell Phone	
Home Address	City □	State, Zip code
Student's Diagnosis: <b>DIABETES:</b> <input type="checkbox"/> Type 1 <input type="checkbox"/> Type 2 <input type="checkbox"/> Other	Today's Date	

MONITORING		
<b>BLOOD GLUCOSE (BG) MONITORING</b> with meter, lancets, lancing device, and test strips	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Student requires supervision <input type="checkbox"/> To be performed by school personnel <input type="checkbox"/> Student is independent <input type="checkbox"/> Permission to self-carry	<input type="checkbox"/> Before meals <input type="checkbox"/> For symptoms of hypo/hyperglycemia & anytime the student does not feel well <input type="checkbox"/> Before PE/Activity <input type="checkbox"/> After PE/Activity <input type="checkbox"/> Prior to dismissal <input type="checkbox"/> Additional BG monitoring may be performed at parent's request
<b>CONTINUOUS GLUCOSE MONITORING (CGM)</b>  Brand/Model: _____	<input type="checkbox"/> Yes <input type="checkbox"/> No Alarms set for: Low: _____ (mg/dL) High: _____ (mg/dL)	Always confirm CGM results with finger stick check before taking action on sensor blood glucose level. If student has symptoms or signs of hypoglycemia, check finger stick blood glucose level regardless of CGM.
<input type="checkbox"/> <b>URINE KETONE TESTING</b> <input type="checkbox"/> <b>BLOOD KETONE TESTING</b>	Anytime the BG > _____ mg/dL or when student complains of nausea, vomiting, abdominal pain. See page 3 for further instructions under hyperglycemia management.	

NAME OF MEDICATION	DOSE/ROUTE		TIME	
<input checked="" type="checkbox"/> <b>GLUCAGON</b> - INJECTABLE	<input type="checkbox"/> 0.5 mg subq/IM <input type="checkbox"/> 1.0 mg subq/IM		Immediately for severe hypoglycemia: unconscious, semi-conscious (unable to control his/her airway or unable to swallow), or seizing	
	<b>DOSAGE</b>	<b>TIME</b>	<b>POSSIBLE SIDE EFFECTS</b>	<b>TREATMENT OF SIDE EFFECTS</b>
<input type="checkbox"/> Glucophage® (Metformin) <input type="checkbox"/> to be administered at school	_____ mg po	_____ AM or PM	Nausea/vomiting, diarrhea	Clear liquids
<input type="checkbox"/> Other: _____® <input type="checkbox"/> to be administered at school				

**Additional Instructions:**

Specific duration of order: <b>2009-2010 SCHOOL YEAR</b>	Physician/Provider Signature: _____ Provider Printed Name: _____ □	Office Phone: _____ Office Fax: _____ <b>Emergency #</b> _____
---	--	--

Institution Name and Address:

DIABETES MEDICAL MANAGEMENT PLAN  
INTENSIVE THERAPY

Page 2 of 3

Patient Label or MRN, Acct#, Patient name, DOB, Date of Service

**SCHOOL YEAR 2009-2010 DIABETES SCHOOL CARE PLAN**

Student: \_\_\_\_\_

**Intensive Therapy/Multiple Daily Injections**

Effective date: \_\_\_\_\_

**Definitions**

<p><b>Insulin-to-Carbohydrate Ratio (CHO Ratio)</b></p> <ul style="list-style-type: none"> <li>the amount of insulin necessary to prevent hyperglycemia after ingestion of a specified amount of carbohydrate</li> <li>usually expressed as "1 unit for every ____ grams of carbohydrate"</li> </ul>	<p><b>Insulin Sensitivity (Correction Factor)</b></p> <ul style="list-style-type: none"> <li>the predicted drop in blood glucose concentration after administration of 1 unit of regular or rapid-acting insulin</li> <li>usually expressed as "1 unit for every ____ mg/dl blood glucose is &gt; target"</li> </ul>	<p><b>Target Blood Glucose</b></p> <ul style="list-style-type: none"> <li>a specific blood glucose value used to determine the correction dose of insulin administered with a meal</li> </ul>
--	--	---

**INSULIN**

Insulin to be given during school hours: <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> May calculate/give own injections with supervision <input type="checkbox"/> Requires assistance to calculate/give injections <input type="checkbox"/> Independently calculates/gives own injection
<input type="checkbox"/> Rapid-acting Insulin Type: _____ <sup>®</sup> <i>(all doses to be administered subcutaneously)</i>	<b>Timing of Insulin Dose:</b> Rapid-acting Insulin should always be given prior to <input type="checkbox"/> meals <input type="checkbox"/> snacks if CHO intake can be predetermined. ➤ If CHO intake cannot be predetermined insulin should be given no more than 30 minutes after completion of meal/snack. ➤ Treat hypoglycemia before administration of meal or snack insulin.	
<input type="checkbox"/> _____ <sup>®</sup> _____ units at _____ am or pm <input type="checkbox"/> may mix with rapid-acting insulin <i>(all doses to be administered subcutaneously)</i>	<b>CALCULATING INSULIN DOSES:</b> According to CHO ratio and Insulin Sensitivity/Correction Factor (if needed) - the student requires meal time coverage with rapid-acting insulin based on the amount of carbohydrates in the meal and may require additional insulin to correct blood glucose to the desired range according to the following formula:  <b>Insulin Dose = [(Actual BG – Target pre-meal BG) divided by Insulin Sensitivity] + [# carbohydrates consumed/CHO Ratio]</b>  <ul style="list-style-type: none"> <li>Fractional amounts of insulin from correction and carbohydrate calculation, when added together, may yield an even amount of insulin</li> <li>If uneven, then round to the nearest <b>half or whole unit</b> (May use clinical discretion; if physical activity follows meal, then may round down).</li> </ul>	
Target pre-meal BG: _____ mg/dL	<b>Insulin Sensitivity/Correction Factor:</b> <input type="checkbox"/> unit for every _____ > target	
<b>CHO Ratio:</b> <input type="checkbox"/> Parent has permission to adjust CHO ratio in a range from 1:_____ to 1:_____	<b>Exercise/PE CHO Ratio:</b> _____ <input type="checkbox"/> <b>Not Applicable</b> <ul style="list-style-type: none"> <li>Less insulin may be required with meals prior to physical activity in order to prevent hypoglycemia. If so, the Exercise/PE CHO Ratio should be used instead of the CHO Ratio.</li> </ul>	
<input type="checkbox"/> Correction insulin to be administered for elevated blood glucose if 3 hours or more after last insulin dose		

**Snacks**

- In general, children with diabetes managed using Intensive Therapy/MDI do not require snacks.
- Scheduled snacks may be required prior to or after exercise in order to prevent hypoglycemia. Insulin is not administered with these snacks.  
 Before Exercise  After Exercise
- Foods may be eaten at unscheduled times. Insulin may be ordered for these snacks in order to prevent post-meal hyperglycemia (see above).
- Snack time insulin = # carbohydrates consumed/CHO Ratio.
- Never provide insulin coverage for carbohydrate/glucose being used to treat hypoglycemia.

**Exercise and Sports**

- In general, there are no restrictions on activity unless specifically noted.
- A student should not exercise if his/her blood glucose is < 70 mg/dL or > 300 mg/dL (with positive ketones) immediately prior to exercise or until hypoglycemia/hyperglycemia is resolved.
- A source of fast-acting glucose & glucagon should be available in case of hypoglycemia.

Specific duration of order: <b>2009-2010 SCHOOL YEAR</b>	Physician/Provider Signature: _____ <input type="checkbox"/>	Provider Printed Name: _____	Office Phone: _____ Office Fax: _____ <b>Emergency #</b> _____
---	---	------------------------------	--

Institution Name and Address:

**DIABETES MEDICAL MANAGEMENT PLAN  
INTENSIVE THERAPY**

Page 3 of 3

Patient Label or MRN, Acct#, Patient name, DOB, Date of Service

**SCHOOL YEAR 2009-2010 DIABETES SCHOOL CARE PLAN**

**Student:** \_\_\_\_\_

**Effective date:** \_\_\_\_\_

**Hypoglycemia (Low Blood Glucose)**

Hypoglycemia is defined as a blood glucose  $\leq$  \_\_\_\_\_ mg/dL

Signs of hypoglycemia:

Hunger	Sweating	Shakiness	Paleness	Dizziness
Confusion	Loss of coordination	Fatigue	Fighting	Crying
Day-dreaming	Inability to concentrate	Anger	Passing-out	Seizure

- If hypoglycemia is suspected, check the blood glucose level.

<b>Hypoglycemia Management (Low Blood Glucose)</b>	<b>Severe Hypoglycemia: If student unconscious, semi-conscious (unable to control his/her airway or unable to swallow) or seizing, administer glucagon.</b>
	<ul style="list-style-type: none"> <li>• Place student in the "recovery position."</li> <li>• If glucagon is administered, call 911 for emergency assistance, and call Parents/Legal Guardian.</li> </ul>
	<b>Mild or Moderate Hypoglycemia: If conscious &amp; able to swallow, immediately give 15 grams fast-acting glucose:</b>
	<ul style="list-style-type: none"> <li>• 3-4 glucose tablets or</li> <li>• 6 Life Saver® Candies or</li> <li>• 4 ounces of regular soda/juice or</li> <li>• 1 small tube Glucose/Cake gel</li> </ul>
	<b>Repeat BG check in 15 minutes</b>
	<ul style="list-style-type: none"> <li>• If BG still low, then re-treat with 15 gram CHO</li> <li>• If BG in acceptable range and at lunch or snack time, let student eat and cover CHO per orders</li> <li>• If BG in acceptable range and not lunch or snack time, provide student slowly-released CHO snack (Example: 3-4 peanut butter or cheese crackers or ½ sandwich)</li> </ul>
	If unable to raise the BG > 70 mg/dL despite fast-acting glucose sources, call: _____

**Hyperglycemia (High Blood Glucose)**

Signs of hyperglycemia:

Extreme thirst	Frequent urination	Blurry Vision	Hunger	Headache
Nausea	Hyperactivity	Dry Skin	Dizziness	Stomach ache

- If hyperglycemia is suspected, check the blood glucose level.

<b>Hyperglycemia Management (High Blood Glucose)</b>	<b>If BG &gt; ____ mg/dL, or when child complains of nausea, vomiting, and/or abdominal pain, ask the student to check his/her urine for ketones</b>
	<ul style="list-style-type: none"> <li>• If urine ketones are trace to small (blood ketones 0 - 1.0 mmol/L), give 8-16 ounces of sugar-free fluid (water), return to classroom</li> <li>• If correction insulin has not been administered within 3 hours, provide correction insulin according to student's Correction Factor and Target pre-meal BG</li> <li>• Recheck BG and ketones 2 hours after administering insulin</li> </ul>
	<ul style="list-style-type: none"> <li>• If urine ketones are moderate/large (blood ketones &gt;1.0 mmol/L), give 8-16 ounces of sugar-free fluid (water) and call _____ for instructions concerning insulin administration.</li> <li>• Contact the Parent/Legal Guardian.</li> <li>• Recheck BG and ketones 2 hours after administering insulin</li> </ul>

My signature below provides authorization for the above written orders. I/We understand that all treatments and procedures may be performed by the school nurse, the student and / or trained unlicensed designated school personnel under the training and supervision provided by the school nurse (or by EMS in the event of loss of consciousness or seizure) in accordance with state laws & regulations. I also give permission for the school to contact the health care provider regarding these orders and administration of these medications.

School plan ordered by:	Physician/Provider Signature: _____	Provider Printed Name: _____	Date: _____
Acknowledged and received by:	Parent/Legal Guardian: _____		Date: _____
Acknowledged and received by:	School Representative: _____		Date: _____

Institution Form #